

Antipsychotics Prescribed

I4C Indicator MED-PSY-002: Antipsychotics Prescribed v 1.1

Category: Medication Management/Psychoactive

Purpose:

This indicator is used to identify patients prescribed an antipsychotic medication.

Source:

Based on Health Data Coalition Current Clinical Measures, June 2018.

Base Population:

All patients with an Active demographic status recorded in the EMR.

Indicator Segments:

Currently prescribed: Count of patients WITH at least one current antipsychotic prescription documented.

Only past prescriptions: Count of patients WITHOUT ANY current antipsychotic prescriptions documented AND WITH at least one past antipsychotic prescription documented.

Never Prescribed: Count of patients WITHOUT ANY current or past antipsychotic prescriptions documented.

Current antipsychotic prescription is any of the following medications (generic names listed) that are current (not discontinued) medications:

'aripiprazole'	'asenapine'	'brexpiprazole'	'clozapine'
'lurasidone'	'olanzapine'	'paliperidone'	'quetiapine'
'risperidone'	'ziprasidone'	'loxapine'	'pimozide'
'fluspirilene'	'chlorpromazine'	'fluphenazine'	'levomepromazine'
'mesoridazine'	'periciazine'	'perphenazine'	'pipotiazine'
'prochlorperazine'	'promazine'	'thioridazine'	'trifluoperazine'
'thiopropazine'	'thiothixene'	'flupentixol'	'tiotixene'
'haloperidol'	'zuclopenthixol'		

Past antipsychotic prescription is any of the above medications that are past (discontinued) medications.

Suggested Indicator Use:

Physician or Practice use of this indicator is to identify or recall patients who are currently prescribed antipsychotic medications.